

- mm. Postpetiole height 0.23 mm. Known from Kazakhstan and Kirghizia (Fig. 268) *S. picetojuglandeti*
Figs. 20, 21, 57, 63, 79, 91, 251 - 255.
- b. Gastral tergite one with basal carinae as long as $\frac{1}{2}$ length of postpetiole. Remaining combination of characters not matching those listed above (Fig. 327) *S. ussuriense*
Figs. 61, 65, 83, 95, 310 - 314.
- 16 (5) a. Mandible with 11 teeth. Compound eye length always less than 0.10 mm. Gastral tergite one with basal carinae as long as $\frac{1}{2}$ length of postpetiole. Metanotal impression deep; depth approximately $\frac{1}{2}$ length of propodeal spines. Known from Japan (Fig. 209)
..... *S. nipponense*
Figs. 52, 70, 86, 200 - 202.
- b. Mandible usually with 9 or fewer teeth. Remaining combination of characters not matching those listed above 17
- 17 (16) a. Mandible with 6 teeth. Propodeal plates shorter than propodeal spines. Gastral tergite one with basal carinae as long as $\frac{1}{3}$ length of postpetiole or shorter. Gastral sternite one lacking basal carinae and glassy-smooth. Propodeal plates approximately 0.75 length of propodeal spines. Thoracic dorsum rugose. Known only from type series, Nepal (Fig. 130) *S. gurkhalis*
Figs. 51, 74, 87, 126 - 129.
- b. Mandible with 7 or more teeth. Remaining combination of characters not matching those listed above 18
- 18 (17) a. Ocular index 14% or greater 19
- b. Ocular index typically less than 12% 21
- 19 (18) a. Gastral tergite one with basal carinae as long as $\frac{1}{3}$ to $\frac{1}{2}$ length of postpetiole. Gastral sternite one with basal carinae as long as 0.20 to 0.25 length of postpetiole (Fig. 110) *S. debile*
Figs. 10 - 17, 32 - 34, 36 - 39, 44, 45, 47, 100 - 102.
- b. Gastral tergite one with basal carinae as long as $\frac{1}{4}$ to 0.20 length of postpetiole. Gastral sternite one lacking basal carinae and glassy-smooth 20
- 20 (19) a. Compound eye length 0.11 - 0.18 mm. Postpetiole length 0.24 - 0.40 mm. Postpetiole width 0.21 - 0.29 mm. Known from Japan and China (Fig. 236) *S. owstoni*
Figs. 49, 71, 84, 226 - 228.
- b. Compound eye length 0.08 - 0.11 mm. Postpetiole length 0.21 - 0.26 mm. Postpetiole width 0.19 - 0.23 mm. Known only from type series, Pakistan (Fig. 153) *S. jeritorum*
Figs. 53, 73, 148 - 152.
- 21 (18) a. Gastral sternite one lacking basal carinae and glassy smooth