



FIG. 1—1) Lateral view of mesosoma, petiole (p) and postpetiole (pp); and 2) dorsal view of petiole (p) and postpetiole (pp) of holotype worker of *N. isodontatus*; 3) mandible (right) of larger worker paratype of *N. isodontatus*; 4) mandible (left) of holotype of *N. isodontatus*; and 5) head (full face view) of the holotype worker of *N. isodontatus*. All figures are drawn to the same scale.

mesonotum and pronotum (Fig. 1-1), broadly angulate between dorsal face and posterior face, but without transverse carina, dorsal face and posterior face about equal in length, posterior face not concave and without carinae along lateral edges; petiole node about as wide as long, wider posteriorly (Fig. 1-2); postpetiole wider than long.

Most body surfaces with hairs about 0.1 mm in length, anterior border of clypeus with a row of dense hairs (Fig. 1-5); funiculus with dense, short, bristly hairs; decumbent pubescence absent. Body surfaces are polished and clear

golden yellow in color. Large punctures are present on most body surfaces, and are most evident on the head and dorsum of mesosoma.

*Female and Male*—Unknown.

*Type Series*—Holotype worker (Museum of Comparative Zoology, Harvard University) and 24 paratype workers (United States National Museum, Texas A&M University, British Museum of Natural History, Museo de Historia Nacional, Colombia, Museu de Zoologia da Universidade de São Paulo, collection of Dr. Julian Watkins, and the Laboratory for Environmental Biology, University of Texas); USA, Texas,