

- Ali, S. M.
1959. Female reproductive system of *Pseudococcus vastator* Maskell (Hemiptera: Pseudococcidae). Indian J. Entomol. 20: 142–146.
- Balachowsky, A. S.
1957. Sur un nouveau genre aberrant de cochenille radicicole myrmecophile nuisible au cafeier en Colombie. Rev. Path. Veg. Entomol. Agric France. 36: 157–164.
- Beardsley, J. W.
1969. A new fossil scale insect (Homoptera: Coccoidea) from Canadian amber. Psyche 76: 270–279.
1970. Three new species of *Chavesia* Balachowsky from Tropical America (Homoptera: Coccidae). Proc. Hawaiian Entomol. Soc. 20(3): 509–520.
- Borgmeier, J. W.
1927. Um cave de trophobiose entre uma formiga e um parasito de cafeiro. Boll. Mus. Nac. Rio Janeiro 3: 285–289.
- Boucot, A. J.
1990. Evolutionary paleobiology of behavior and coevolution. Amsterdam: Elsevier.
- Brandão, C. R. F., R. G. Martins-Neto, and M. A. Vulcano
1990. The earliest known fossil ant (first southern hemisphere mesozoic record) (Hymenoptera: Formicidae: Myrmicinae). Psyche 96: 195–208.
- Brown, W. L., Jr.
1945. An unusual behavior pattern observed in a Szechuanese ant. J. West China Border Res. Soc. B 15: 185–186.
- Bünzli, G. H.
1935. Untersuchungen über coccidophile Ameisen aus den Kaffeefeldern von Surinam. Mitt. Schweiz. Entomol. Ges. 16: 453–593.
- Buschinger, A., J. Heinze, K. Jessen, P. Douwes, and U. Winter
1987. First European record of a queen ant carrying a mealybug during her mating flight. Naturwiss. 74: 139–140.
- Campos, L. A. de O., and H. C. Morais
1986. Transporte de homópteros por fêmeas de formigas do gênero *Acropyga* (Formicinae) em Viçosa, MG. In F. H. Caetano (ed.), Anais Do VII Encontro de Mirmecologia do Estado de São Paulo, UNESP-FAPESP-CNPq: 52–53.
- Carpenter, F. M.
1992. Superclass Hexapoda, vol. 4 (Part R, Arthropoda 4), Treatise on Invertebrate Paleontology. Lawrence, KS: Geological Soc. America/ U. Kansas.
- Delabie, J. N. C., J. E. Mantovani, and I. Mauricio
1991. Observaçoes sobre a biologia de duas espécies de *Acropyga* (Formicidae, Formicinae, Plagiolepidini) associadas a rizosfera do cacau. Rev. Brasil. Biol. 51: 185–192.
- De Lotto, G.
1977. On some African mealybugs (Homoptera: Coccoidea: Pseudococcidae). J. Entomol. Soc. South Afr. 40: 13–36.
- DeVries, P. J.
1991. Mutualism between *Thisbe irenea* butterflies and ants, and the role of ant ecology in the evolution of larval-ant associations. Biol. J. Linn. Soc. 43: 179–195.
- Dill, M., and U. Maschwitz
1998. The nomadic herdsmen of Kinabalu Park: a unique lifestyle in ants. Sabah Parks Nat. J. 1: 61–67.
- Dlussky, G. M.
1975. Formicoidea, Formicidae, Sphaecomyrminae. In A. P. Rasnitsyn (ed.), The higher Hymenoptera of the Mesozoic. Trans. Paleont. Inst. Acad. Sci. USSR 147: 115–122. [In Russian]
1983. A new family of Upper Cretaceous Hymenoptera—an “intermediate link” between the ants and the scolioids. Paleontol. Zh. 3: 65–78.
1987. New Formicoidea (Hymenoptera) of the Upper Cretaceous. Paleontol. J. 1987: 146–150.
1988. Ants from (Palaeocene?) Sakhalin amber. Paleontol. J. 1: 50–61. [English translation of original Russian version by Scripta Technica]
- Eberhard, W. G.
1978. Mating swarms of a South American *Acropygia* (Hymenoptera: Formicidae). Entomol. News 89: 14–16.
- Emery, C.
1925. Hymenoptera, Family Formicidae. In P. Witsman (ed.), Genera Insectorum 183: 1–302.
- Encarnacao, A. V. Da, J. N. C. Delabie, and D. Matile-Ferrero
1993. Novas observaçoes sobre a biologia de *Acropyga* s. pp. e dos seus Pseudococcidae associados nos cacauais da Bahia (Hymenoptera: Formicidae e Homoptera: Coccoidea). Anais da 45a reunião Anual da SBPC, Vol. 1: 931. U.F. Pernambuco, Recife, Brazil.
- Fiedler, K., and U. Maschwitz
1989. Functional analysis of the myrmecophilous relationships between ants (Hym.: