

Kashmir, 4 km S Kulan, 34°14' N, 75°10' E, 3599 m, No. 3046, 3048, 07.08.1978; 3 workers, [India], Kashmir, Sain Nar, 34°06' N, 75°34' E, 3750 m, No. 3054, 11.08.1978; 4 workers, 1 female, 4 males, [India], Kashmir, Wampet, 34°04' N, 75°37' E, 3700 m, No. 3057, 12.08.1978; 1 male, [India], Kashmir, Up. Kaintal Nar, Oitto, 34°00' N, 75°45' E, 4200 m, No. 3068 (all leg. P. Ward); 3 workers, India, Beastel, 20 km S Rhotang, 2800 m, No. J-23, 18.08.1990, leg. J. Heinze (NHML, NHMB, RIGATO, SCHULZ, WARD, SIZK, ELMES, MEI).

Workers. Head long with parallel sides, straight or very feebly convex occipital margins, and narrowly rounded occipital corners. Anterior clypeal margin prominent

Table 5. The mean, minimum and maximum indices of males and queens of six new *Myrmica* species. The measurements codes are as indicated in the methods and the number of individuals measured are given in parenthesis

Табл. 5. Среднее, минимальное и максимальное значение индексов для самок и самцов 6 видов *Myrmica*, подсчитанные как указано в тексте; количество измеряемых экземпляров приведено в скобках

	♀ <i>M. villosa</i>	♀ <i>M. nitida</i> (3)		♀ <i>M. wardi</i> (2)		♀ <i>M. ordinaria</i>	♀ <i>M. rhytida</i> (16)	
		Mean	Min-Max	Mean	Min-Max		Mean	Min-Max
CI	1.15	1.20	1.19-1.20	1.20	1.16-1.24	1.27	1.20	1.17-1.24
FI	0.46	0.45	0.44-0.46	0.41	0.40-0.43	0.47	0.42	0.41-0.43
FLI	1.06	1.03	1.03	1.035	1.03-1.04	1.07	1.05	1.02-1.07
SI1	0.87	0.75	0.74-0.76	0.76	0.73-0.80	0.74	0.75	0.71-0.80
SI2	1.00	0.90	0.88-0.91	0.92	0.90-0.93	0.93	0.90	0.86-0.96
PI1	1.28	1.22	1.19-1.25	1.37	1.25-1.48	1.54	1.43	1.27-1.56
PI2	0.47	0.50	0.49-0.50	0.51	0.48-0.55	0.60	0.57	0.51-0.61
PPI1	0.91	0.83	0.80-0.85	0.98	0.96-1.00	0.76	0.90	0.83-0.96
PPI2	0.94	1.01	1.00-1.02	0.94	0.91-0.97	1.05	0.97	0.91-1.00
PPI3	1.48	1.52	1.50-1.52	1.64	1.61-1.67	1.69	1.55	1.48-1.64
ESLI	0.29	0.20	0.18-0.20	0.23	0.21-0.25	0.27	0.20	0.17-0.25
ESDI	1.74	2.33	2.21-2.54	2.04	1.96-2.11	1.67	2.13	1.88-2.50
HTI	0.94	0.96	0.96-0.96	0.96	0.95-0.96	0.98	0.90	0.83-0.97
AI	1.90	1.68	1.63-1.72	1.65	1.63-1.67	1.74	1.69	1.62-1.75
SCI	1.53	1.45	1.43-1.48	1.50	1.50-1.50	1.52	1.47	1.28-1.59
WI1							6.62	6.43-6.79

Table 5 (cont.)

Таблица 5 (продолжение)

	♀ <i>M. petita</i>	♂ <i>M. nitida</i>	♂ <i>M. wardi</i> (2)		♂ <i>M. rhytida</i> (12)	
			Mean	Min-Max	Mean	Min-Max
CI	1.24	1.11	1.08	1.05-1.11	1.14	1.11-1.16
FI	0.41					
FLI	1.09					
SI1	0.73	0.88	0.52	0.49-0.56	0.90	0.85-0.95
SI2	0.90	0.98	0.56	0.54-0.58	1.03	0.98-1.09
PI1	1.35	1.25	1.34	1.33-1.36	1.41	1.14-1.59
PI2	0.56	0.57	0.64	0.62-0.65	0.62	0.56-0.69
PPI1	0.84	1.00	0.86	0.85-0.87	0.89	0.82-0.98
PPI2	1.00	1.03	1.02	1.00-1.03	0.97	0.88-1.04
PPI3	1.58	1.36	1.40	1.38-1.42	1.48	1.23-1.58
ESLI	0.29					
ESDI	1.67					
HTI	0.90	1.30	1.45	1.43-1.46	1.30	1.21-1.46
AI	1.68	1.49	1.65	1.62-1.68	1.63	1.55-1.68
SCI	1.57	1.33	1.22	1.21-1.23	1.33	1.16-1.23
WI1			7.50	7.46-7.55	6.94	6.27-7.55