

(NC), Bahia Honda (GU, MD), Big Pine (EW, NC, JT, GU, MD), No Name (JT, MD), Sugarloaf (MD), Shark (NC), Stock Island (MD), Dry Tortugas (CW, NC). Vouchers: ABS, NC, JT, GU.

*Solenopsis gobularia littoralis* Creighton. This native southeastern species lives in open sandy areas and can also be found in tussocks in salt marshes. Elliott (MD), Largo (MD), Plantation (EW), Ohio (S&JP), Big Pine (EW, JT, GU, MD), Saddlebunch 2 (JW), Geiger (JW). Vouchers: ABS, JT, GU.

*Solenopsis invicta* Buren. This exotic species is not nearly as dominant in the Keys as elsewhere in Florida, possibly because *S. geminata* is better adapted to coastal areas. Nests are in soil, usually in moist unshaded sites. Elliott (MD), Saddlebunch 5 (NC), Key West (NC). Vouchers: ABS, NC.

*Solenopsis picta* Emery. This native southeastern species nests in twigs, holes in the bark of live pine trees, and plant stems. Elliott (MD), Plantation (EW), Upper Matecumbe (EW), Bahia Honda (MD), Key West (MD). Vouchers: ABS.

*Solenopsis tennesseensis* M. R. Smith. This native southeastern species occurs in soil or leaf litter in wooded areas. Elliott (MD), Largo (EW, MD), Plantation (MD), Grassy (JT, MD), Long (MD), Bahia Honda (MD), No Name (JT, MD), Boca Chica (JW), Stock Island (MD). Vouchers: ABS, JT.

*Solenopsis* sp. This small yellow species in the subgenus *Diphorhoptum* is found throughout Florida. It has not been formally described. This species is probably native to the Keys. Nests are in leaf litter, usually in shaded habitats. Elliott (MD), Largo (MD), Plantation (DS, MD), Lower Matecumbe (MD), Grassy (JT, MD), Bahia Honda (JT, MD), Big Pine (JT, MD), No Name (MD), Sugarloaf (MD), Boca Chica (JW), Stock Island (MD), Key West (JW, MD). Vouchers: ABS, JT.

*Strumigenys eggersi* Emery. This neotropical species is probably exotic in Florida; it is currently found through the southern two-thirds of the state. Nests are in leaf litter in shaded or partially shaded areas. Largo (MD), Plantation (DS, MD), Upper Matecumbe (MD), Big Pine (MD, K&P), Sugarloaf (JW), Stock Island (K&P, MD), Key West (JW). Vouchers: ABS, GU.

*Strumigenys gundlachi* (Roger). This neotropical species is probably exotic in Florida; it is found only in the extreme southern part of the state. Nests are in leaf litter in shaded sites. Elliott (MD), Largo (EW, MD), Upper Matecumbe (MD), Grassy (MD), Long (MD), Vaca (K&P), Big Pine (MD, K&P), No Name (JT, MD), Sugarloaf (MD, K&P). Vouchers: ABS, JT, GU.

*Strumigenys louisianae* Roger. The range of this species extends from the southern U.S. to southern South America. It is presumably native in Florida. Nests are in leaf litter in moist shaded areas. Largo (JT), Plantation (MD), Big Pine (K&P), No Name (MD, GU), Big Torch (S&JP). Vouchers: ABS, JT, GU.

*Strumigenys silvestrii* Emery. This neotropical species is probably exotic in Florida. It is known from a few widely scattered localities in the state. Nests are in leaf litter. Plantation (DS), Big Pine (MD), West Summerland (K&P). Vouchers: ABS, GU.

*Tapinoma litorale* Wheeler. This species occurs in the West Indies and the southern third of Florida. We consider this a native species. Nests are in twigs and vines, usually in exposed or partially exposed sites. Elliott (MD), Largo (EW), Plantation (EW), Windley (EW), Big Pine (EW, GU, MD), Long (MD), Bush (NC). Vouchers: ABS, NC, GU.

*Tapinoma melanocephalum* (Fabricius). This exotic species is common in disturbed habitats through southern Florida. Nests are under bark, at the bases of palm leaves, and other similar cavities. Elliott (MD), Largo (MD), Plantation (DS), Upper Matecumbe (MD), Grassy (MD), Long (MD), Vaca (GU), Big Pine (MD, GU), Saddlebunch 2 (JW), Stock Island (MD), Key West (JW, MD), Dry Tortugas (CW, NC). Vouchers: ABS, NC, GU.